

CAN CLUSTER EVAPORATION EXPLAIN THE MISSING THERMAL ENERGY IN GALAXY CLUSTERS?

MIKHAIL V. MEDVEDEV¹

Department of Physics and Astronomy, University of Kansas, KS 66045

Draft version February 5, 2008

ABSTRACT

Recent observations of a number of galaxy clusters using the Sunyaev-Zel'dovich effect indicate that about 1/3 of baryonic mass is missing from the hot intracluster medium (ICM), which is significantly larger than the fraction of stars and cool gas, which account for only about 10%. Here we address the question whether the remaining $22 \pm 10\%$ can be accounted for by thermal evaporation of gas from clusters. We have found that evaporation can occur only from the cluster “surface”, $r \sim r_{\text{vir}}$, and not from its interior. We evaluated particle diffusion through the magnetized ICM for several scenarios of ISM turbulence and found that diffusivity is suppressed by at least a factor of 100 or more, compared to the Spitzer value. Thus, only particles from radii $r \gtrsim 0.9r_{\text{vir}}$ can evaporate. Diffusion of particles from inside the cluster, $r \lesssim 0.9r_{\text{vir}}$, takes longer than the Hubble time. This lowers the cluster-averaged fraction of the evaporated hot gas to few percent or less. However, if the missing hot component is *indeed* due to evaporation, this strongly constrains the magnetic field structure in the cluster envelope, namely either (i) the gas is completely unmagnetized ($B \leq 10^{-21}$ gauss) in the cluster halo or (ii) the magnetic fields in the ICM are rather homogeneous and non-turbulent.

Subject headings: galaxies: clusters: general — diffusion — magnetic fields

1. INTRODUCTION

Observations of the Sunyaev-Zel'dovich effect in galaxy clusters directly probe the energy content of the intracluster medium (ICM) in the model-independent way, and it is complementary to X-ray observations (see Voit 2005 and references therein). The study of 193 clusters with temperatures above 3 keV using the cosmic microwave background data from WMAP yielded a remarkable and puzzling result (Afshordi, et al. 2006). It has been shown that SZ effect would account for a half of the thermal energy of the ICM provided all the baryons are in the hot phase. This result implies that a significant fraction of baryons, about $32 \pm 10\%$ by mass, is missing from the ICM phase.

It has been proposed (Loeb 2007) that gas evaporation can reduce internal energy of gas in a cluster. Evaporation involves particles from the thermal tail of the Maxwellian distribution because their velocities exceed the escape velocity from a cluster, hence they are gravitationally unbound and can leave the cluster. Since these particles are suprathermal, i.e., their speed exceeds the thermal speed, the overall energy content of the cluster will be decreasing with time. Because the ICM gas is mildly collisional, the transport of these particles will be diffusive. However, in the presence of magnetic fields in the ICM, particle diffusion is often substantially reduced (Rechester & Rosenbluth 1978; Chandran & Cowley 1998; Narayan & Medvedev 2001; Malyshkin & Kulsrud 2001; Malyshkin 2001; Lazarian 2006). The reduction coefficient $f_B < 1$ has been introduced; its value has not been accurately evaluated, however.

Magnetic fields in galaxy clusters are highly turbulent with the fluctuating component being of order the mean field, $\delta B \sim B$, and is tangled on small scales. The magnetic field spectrum deduced from the rotation mea-

sure observations is peaked at about $l_B \sim \text{one} - \text{few kpc}$, which sets the characteristic correlation length of the field (Vogt & Enßlin 2003, 2005; Enßlin, et al 2005). We evaluate the diffusion suppression coefficient $f_B = \kappa_{\text{eff}}/\kappa_{\text{Sp}}$ (where κ_{eff} and κ_{Sp} are the effective and Spitzer diffusion coefficients) in the turbulent magnetized ICM. The obtained value turns out to be about 0.01 for a range of typical parameters of the ICM gas. This value is substantially lower than what have previously been assumed. We discuss implications of this result for the cluster properties, dynamics and formation.

2. PARTICLE DIFFUSION IN THE CLUSTER HALO

For simplicity we assume that the ICM gas is hydrogen and neglect the effect of helium. We also assume that the magnetic field in the ICM is highly turbulent and chaotic. Particle transport in such a system is strongly influenced by the field properties.

A particle in the magnetic field gyrates about a certain field line and can move freely along it. Thus, on average, the particle closely follow the field line. A particle can jump through the distance $\sim r_L = v_{\text{th}} m_p c / eB$ (the Larmor radius) to a neighboring field line in particle-particle collisions. Since the Larmor radius is many orders of magnitude smaller than the mean-free-path in the ICM, the cross-field diffusion is greatly suppressed. For future reference, the Larmor radius, the proton mean-free-path for the typical conditions in the galaxy cluster halo, $r \sim r_{\text{vir}}$, and the virial radius are $r_L = 2.3 \times 10^{-12}$ Mpc $T_{5\text{keV}}^{1/2} B_{\text{lnG}}^{-1}$, $\lambda = 1.3$ Mpc $T_{5\text{keV}}^2 \Delta_{25}^{-1}$, $r_{\text{vir}} = 2.3$ Mpc $T_{5\text{keV}}^{1/2}$, where $\Delta = n_p/\bar{n}_p \sim 25$ is the overdensity at the present epoch, $z = 0$, characterizing the ICM gas density at the virial radius, $\bar{n}_p = 2 \times 10^{-7} \text{ cm}^{-3}$ is the mean density in the Universe at $z = 0$ (Spergel, et al. 2006),

¹also at the Institute for Nuclear Fusion, RRC “Kurchatov Institute”, Moscow 123182, Russia

$\Delta_{25} = \Delta/25$, $T_{5\text{keV}} = T/(5 \text{ keV})$, the ICM magnetic field $B_{1\text{nG}} = B/(10^{-9} \text{ gauss})$. The thermal and Alfvén speeds are $v_{\text{th}} = (3kT/m_p)^{1/2} = 1.2 \times 10^8 T_{5\text{keV}}^{1/2} \text{ cm s}^{-1}$, $v_A = B/(4\pi m_p n_p)^{1/2} = 9.7 \times 10^4 B_{\text{nG}} \Delta_{25}^{-1/2} \text{ cm s}^{-1}$. Below we consider three different scenarios for the particle transport: the case of static chaotic fields, the Alfvénic cascade and transport by large-scale turbulent motions (including the hydrodynamic cascade). We show that transport in Alfvénic cascade Lazarian (2006) is suppressed by some power of the Alfvénic Mach number, whereas the static fields and the fluid motions yield similar values for the diffusion coefficient for typical cluster conditions.

Static chaotic fields — In a medium with strong magnetic turbulence, the magnetic field lines themselves can be chaotic, such that the separation of two field lines increases exponentially with distance along them, $d \sim r_L \exp(l_{\parallel}/l_B)$, with l_B being the field correlation length.

The transport of a proton along such field lines results in effective diffusion coefficient is suppressed compared to the classical Spitzer value $\kappa_{\text{Sp}} = \lambda v_{\text{th}} = 4.8 \times 10^{32} \text{ cm}^2 \text{ s}^{-1} T_{5\text{keV}}^{5/2} \Delta_{25}^{-1}$. Note that here we shall not use the ambipolar rate, which is a factor of two larger.²

If the magnetic field has a well defined, single scale l_B , the suppression factor due to the field line geometry, i.e., the field line tangling alone, is (Rechester & Rosenbluth 1978; Chandran & Cowley 1998) $f_{\text{field}} \simeq [3 \ln(l_B/r_L)]^{-1} \sim 1/60$. Here the factor 1/3 appears because the particle motion is effectively one-dimensional (i.e., along a field line), rather than three-dimensional. Here we also used the observational value of the typical field correlation scale $l_B \sim 1 \text{ kpc}$ (Vogt & Enßlin 2003, 2005).

If l_B ranges through two decades or more in scale, as one can expect in the inertial range of strong magneto-hydrodynamic (MHD) turbulence, the suppression due to tangling is weaker (Narayan & Medvedev 2001):

$$f_{\text{field}} \simeq 1/5. \quad (1)$$

In this case, the velocity on the injection scale is $V_L = v_A$. For a general case $V_L \neq v_A$, see Lazarian (2006).

Another effect that is limiting the diffusion rate is the magnetic mirroring (Chandran & Cowley 1998; Narayan & Medvedev 2001; Malyskin & Kulsrud 2001). Due to the conservation of the adiabatic invariant of a gyrating particle, $\mu = (mv_{\perp}^2/2)/B$, the particle may not be able to penetrate through the region of strong magnetic field, depending on the particle’s pitch angle and the field strength. In clusters, the field inhomogeneities are large, $\delta B \sim B$, therefore the mirroring effect can be significant.

Accurate calculation of the diffusion suppression due to mirroring in various regimes (single and multi-scale) and with various models of turbulence has been done in an excellent work by Malyskin & Kulsrud (2001). The mirroring suppression depends on the ratio λ/l_B , and it is strong when $\lambda \gg l_B$, as in the cluster halo. Note that the mirroring effect is small in the cluster core where $\lambda < l_B$, hence Narayan & Medvedev (2001) argued that $f_{\text{mirroring}} \lesssim 1$. The results obtained by Malyskin & Kulsrud (2001) can

be approximated quite well as follows:

$$f_{\text{mirroring}} \simeq \min \left\{ [l_B/(10^2 \lambda)]^{1/3}, 1 \right\}. \quad (2)$$

The overall diffusion suppression factor is

$$f_B = f_{\text{field}} f_{\text{mirroring}} \sim 4.3 \times 10^{-3}. \quad (3)$$

Here we used $l_B \sim 1 \text{ kpc}$, and we assumed the most favorable case of multi-scale turbulence with $f_{\text{field}} \sim 1/5$. Note that even if the magnetic fields are correlated on a cluster size scale, $l_B \sim 1 \text{ Mpc}$, the suppression is still rather strong: $f_B \sim 0.05$. Thus, the magnetic field suppression is much stronger than $f_B \sim 1/3$ assumed in (Loeb 2007), for which one have obtained that about 10% of the cluster gas will evaporate during the Hubble time. Simply re-scaling this result to the value of $f_B \sim 4 \times 10^{-3}$ yields the evaporated fraction of order $\sim 0.1\%$. This result is essentially independent of the fraction of open field lines crossing r_{vir} .

Alfvénic turbulence — The particle diffusivity in the Alfvénic MHD cascade (but neglecting the mirroring effect) has been studied by Lazarian (2006). He has found that to sub-Alfvénic turbulence, $v_{\text{gas}}/v_A \equiv M_A < 1$, the diffusion coefficient is suppressed by a factor of $M_A^2 < 1$ compared to the Narayan & Medvedev (2001) result:

$$f_{\text{field}} \sim (1/3)M_A^4, \quad M_A < 1, \quad (4)$$

cf. Eq. (8) in Lazarian (2006, 2007). The factor 1/3 above is approximate, however it is not very different from the factor 1/5 obtained by Narayan & Medvedev (2001).

For the super-Alfvénic turbulence, i.e., gas motions with the Alfvén Mach number $M_A > 1$, there are two regimes. In the first one, the particle mean free path is large compared to the so-called “Alfvén length”, $\lambda > l_A$, where $l_A \approx l_B M_A^{-3} < l_B$, and the suppression is

$$f_{\text{field}} \sim (1/3)(l_A/\lambda) \sim (l_B/\lambda)M_A^{-3}, \quad M_A > 1, \quad (5)$$

cf. Eq. (4) of Lazarian (2006), and note that $l_B \ll \lambda$. In the second case, $\lambda < l_A < l_B$, one recovers the $f_{\text{field}} \sim 1/3$ limit of one-dimensional diffusion. This case, however is not very relevant for cluster evaporation, because the mean free path is comparable to the system size and is much larger than the outer scale of turbulence, $\sim 1 \text{ Mpc} \gg l_B$.

As in any magnetic turbulence, the conservation of the magnetic moment results in additional suppression due to particle trapping, as discussed above, Eqs. (2,3). Since the “field” part of the suppression factor alone is diminished by some power of the Alfvénic Mach number in both sub- and super-Alfvénic regimes, we conclude that the static field model puts less stringent limits on thermal transport.

Fluid turbulence — Turbulent fluid eddies can also contribute to the particle transport. The dynamical diffusion coefficient has been also evaluated by Lazarian (2006):

$$\kappa_{\text{dyn}} \approx \begin{cases} C_d LV_L, & M_A \gtrsim 1; \\ \beta C_d LV_L M_A^3, & M_A < 1, \end{cases} \quad (6)$$

where $\beta \sim 4$ and $C_d \approx 1/3$ (Lesieur 1990) (cf., it is not 2/3 as in Lazarian 2006 because we consider proton transport only). Note that transport in sub-Alfvénic turbulence is

² The ambipolar diffusion coefficient, $\kappa_{\text{amb}} = (T_e + T_p)\kappa_{\text{Sp},e}\kappa_{\text{Sp},p}/(T_p\kappa_{\text{Sp},e} + T_e\kappa_{\text{Sp},p})$, where $\kappa_{\text{Sp},e} = \lambda_e v_{\text{th},e}$ and $\kappa_{\text{Sp},p} = \lambda_p v_{\text{th},p} \ll \kappa_{\text{Sp},e}$ for $T_e \simeq T_p$, following from plasma quasi-neutrality, is commonly used for two-component e^-p plasma. Here, we consider diffusion of a supra-thermal component of the proton distribution through gas, not the global escape of the plasma bulk, hence the Spitzer diffusivity is used.

suppressed by M_A^3 . Here L is the eddy scale and V_L is the turbulent velocity on this scale.

It is instructive to represent the Spitzer diffusivity as

$$\kappa_{\text{Sp}} \approx 0.57 T_{5\text{keV}}^{3/2} \Delta_{25}^{-1} (r_{\text{vir}} v_{\text{th}}), \quad (7)$$

i.e., it is very close to $r_{\text{vir}} v_{\text{th}}$ for typical cluster halo conditions. In the fluid turbulence, the combination LV_L must be substantially smaller than $r_{\text{vir}} v_{\text{th}}$. Indeed, if $LV_L \sim r_{\text{vir}} v_{\text{th}}$, eddies of $L \sim r_{\text{vir}}/2$ and $V_L \sim 2v_{\text{th}}$ break the cluster apart: in virial equilibrium $v_{\text{esc}}^2 = 2v_{\text{th}}^2$ and gas parcels with $V_L > 2^{1/2}v_{\text{th}}$ are gravitationally unbound. Some fraction of gas can be lost in such violent events (e.g., mergers), but this cannot be called a steady-state evaporation.

Numerical simulations of turbulence in clusters, taking into account decaying turbulence, mergers and wakes (Dolag, et al. 2005; Subramanian, et al. 2006; Vazza, et al. 2006) yield a typical value of $LV_L \sim 4.5 \times 10^4 \text{ kpc km s}^{-1} \sim 0.016 T_{5\text{keV}}^{-1} (r_{\text{vir}} v_{\text{th}})$. This yields

$$\kappa_{\text{dyn}}/\kappa_{\text{Sp}} \simeq 0.009 T_{5\text{keV}}^{-5/2} \Delta_{25}, \quad (8)$$

for $M_A > 1$, which is comparable to diffusion in static fields. Whether and how much the ICM fields will limit particle escape through r_{vir} has never been studied, but it goes beyond the diffusion approximation and may require a numerical approach.

Diffusion length and evaporated gas fraction — Considering several scenarios, we found that particle diffusion in the magnetized ICM is suppressed by at least a factor of a hundred or more, compared to the Spitzer value. Thus, the static field scenario represents an optimistic estimate (lowest suppression) and we used it for further analysis. We now evaluate the diffusion length, through which a proton can travel during the Hubble time, $t_H = H^{-1}|_{z=0} \simeq 4 \times 10^{17} \text{ s}$:

$$l_{\text{diff}} = (\kappa_{\text{eff}} t_H)^{1/2} \simeq 0.27 \text{ Mpc } f_{B,0.004}^{1/2} T_{5\text{keV}}^{5/4} \Delta_{25}^{-1/2}, \quad (9)$$

where $f_{B,0.004} = f_B/0.004$. Note that the diffusion coefficient is inversely proportional to the local gas density, which, in turn, is $\propto r^{-3}$ for the NFW profile (Navarro, et al. 1997). Formally replacing Δ with $\Delta(r/r_{\text{vir}})^{-3}$, we see that the diffusion from the inner parts of the cluster is strongly suppressed. Hence, the highly collisional gas at $r < 0.5r_{\text{vir}}$ cannot replenish the evaporated suprathermal component of the particle distribution. Because the escaping supra-thermal protons move at speeds $\gtrsim v_{\text{esc}} \sim \eta^{1/2}v_{\text{th}} \sim 2^{1/2}v_{\text{th}}$, the diffusion distance above is accurate to a factor of order unity. Since most of plasma particles, both thermal and supra-thermal, are effectively trapped, their residence time in between magnetic mirrors can be comparable to the Hubble time. Supra-thermal particles with velocity $v = \eta^{1/2}v_{\text{th}}$ can experience N collisions per the Hubble time: $N \sim t_H/\tau_{\text{coll}} \sim 12 (\eta T_{5\text{keV}})^{-3/2} \Delta_{25}$, where $\tau_{\text{coll}} = \lambda(v)/v = 3.3 \times 10^{16} \text{ s } (\eta T_{5\text{keV}})^{3/2} \Delta_{25}^{-1}$ is the particle collision time, $\eta \sim 2.3$ for v being equal to the escape velocity, v_{esc} , at r_{vir} (Loeb 2007). Thus, there can be significant re-distribution of trapped, $v < v_{\text{esc}}$, and untrapped, $v > v_{\text{esc}}$, components (i.e., an initially trapped supra-thermal proton may become untrapped and vice versa,

due to collisions). This may change, perhaps lower, the effective η . In this regime, numerical computation of the effective η using a self-consistent proton distribution accounting for diffusion, trapping and leaking of particles is highly desirable.

The diffusion length is small fraction of the cluster size:

$$l_{\text{diff}}/r_{\text{vir}} \equiv \xi = 0.12 f_{B,0.004}^{1/2} T_{5\text{keV}}^{3/4} \Delta_{25}^{-1/2}. \quad (10)$$

Thus, cluster evaporation effectively occurs from a thin layer of thickness $\sim l_{\text{diff}} \sim 0.1r_{\text{vir}}$. Particles from radii $r < (r_{\text{vir}} - l_{\text{diff}})$ cannot escape from the cluster within the Hubble time.

The NFW density profile (Navarro, et al. 1997) is $n_{\text{gas}}(x) = n_0/[x(1+x)^2]$, where $x = r/r_s$, $r_s = r_{\text{vir}}/c$, c is the concentration parameter, n_0 is the normalization such that $n(x_{\text{vir}}) = \bar{n}_p \Delta$ and $x_{\text{vir}} = r_{\text{vir}}/r_s = c$. The mass profile of the NFW cluster is $M(x) = M_0[\ln(1+x) - x/(1+x)]$ with $M_0 = 4\pi r_s^3 n_0 m_p$. The fraction of the gas mass in the shell of thickness l_{diff} to the total mass of gas in the entire cluster is

$$\frac{M_{\text{shell}}}{M_{\text{cluster}}} = \frac{\ln\left(\frac{c+1}{c+1-\xi c}\right) - \frac{\xi c}{(c+1)(c+1-\xi c)}}{\ln(c+1) - c/(c+1)}. \quad (11)$$

The supra-thermal fraction of the proton distribution depends on the concentration parameter. It has been evaluated to be $0.064 < \epsilon < 0.092$ at $r = r_{\text{vir}}$ and for $2 < c < 8$ (Loeb 2007). Because of trapping of particles in magnetic traps, the particle distribution can evolve, therefore the actual value of ϵ can differ. The fraction the evaporated gas is, therefore,

$$\mu \equiv \epsilon \frac{M_{\text{shell}}}{M_{\text{cluster}}} \simeq \frac{\epsilon \xi c^2 / (c+1)^2}{\ln(c+1) - c/(c+1)} \sim 7.2 \times 10^{-3}, \quad (12)$$

where we used that $\xi = 0.12$, $c = 4$ and the corresponding $\epsilon = 0.076$. Thus, the evaporation fraction of the ICM gas is about 1%. Figures 1 and 2 represents the evaporated fraction μ as a function of the ICM temperature and the field correlation length, l_B , for several values of c and l_B .

3. DISCUSSION

We evaluated the suppression factor of the diffusive transport of protons in the ICM with magnetic fields in three scenarios: (i) static chaotic fields (Narayan & Medvedev 2001), (ii) Alfvénic and (iii) fluid turbulence (Lazarian 2006). Note that magnetic fields are still required even in case (iii), otherwise the fluid approximation breaks down, because $\lambda \sim r_{\text{vir}}$. We considered diffusion suppression is due to (i) a purely geometric effect of turbulence and (ii) the dynamic effect of particle mirroring and trapping in inhomogeneous fields. We have found that particle diffusion is suppressed to at least 1% of the Spitzer value, and usually much more. For further analysis we used the suppression in static fields as an optimistic representative value.

The smallness of the effective diffusivity, compared to the Spitzer value substantially lowers the overall evaporation gas fraction. We calculated the diffusion distance for typical cluster parameters and found that a particle can diffuse through the distance of about $l_{\text{diff}} \sim 0.1r_{\text{vir}}$ within the Hubble time. Thus, particles that are at radii $r < (r_{\text{vir}} - l_{\text{diff}}) \sim 0.9r_{\text{vir}}$ can never pass through the virial

radius and leave the cluster. Consequently, cluster evaporation occurs from the thin cluster “skin” of thickness $\sim 0.1r_{\text{vir}} \sim 200$ kpc. The fraction of the gas evaporated from this skin can be rather high, perhaps about 10% or even more, comparable to the estimate by Loeb (2007). The exact value shall be obtained from self-consistent simulations taking into account particle trapping, particle collisions, evolution of the distribution function and particle leaking out of the cluster.

We make a prediction that the thermal energy deficit shall be the strongest at the outskirts of the cluster $r \sim r_{\text{vir}}$, in the shell of few hundred kpc in thickness. Deeper in the cluster, the thermal energy deficit shall be negligible. Of course, at radii of few tens kpc, a cooling flow, if present, may lower the thermal energy as well.

Since the diffusion distance is small compared to the cluster size, the fraction of the evaporated gas compared to the total cluster mass is small too. We evaluated it to be less than 1% for typical cluster conditions, $T \sim 5$ keV and $l_B \sim 1$ kpc. Although the concentration parameter c can vary for different clusters, its effect on the evaporated fraction, μ , is not very significant. The ICM gas temperature and the field correlation scale do affect μ . Figures 1,2 show how μ depends on T and l_B . Interestingly, by boosting l_B to ~ 10 kpc and T to ~ 15 keV altogether, one can increase the evaporated fraction to about 2% only. One cannot exceed the value of $\mu \sim \text{few } \%$ even by pushing l_B to the maximum possible value of ~ 3 Mpc, — the cluster

size. Apparently, the predicted evaporated gas fraction is below the observational value of $22 \pm 10\%$ by a large margin.

Now, let’s look at the result from a different point of view. Let’s suppose that the observed thermal energy deficit of $\sim 20\%$ is due to cluster evaporation. What assumptions shall be relaxed in this case? We identify two alternatives. First, the ICM shall be effectively unmagnetized, then the result of Loeb (2007) for the low-suppression scenario holds and the evaporated fraction can be of order 10%. In order to neglect the effects of magnetic field, the particle Larmor radius shall be comparable to the size of a cluster, $r_L \sim r_{\text{vir}}$. This sets the limit on the field strength: $B \leq 10^{-21}$ gauss. Second, if the ICM fields are present, they shall be relatively homogeneous with $\delta B \ll B$. In this case, particle trapping is small and protons can easily escape along open field lines. The evaporated fraction will be reduced by the fraction of the cluster surface threaded by open fields lines. If this suppression is not strong, i.e., of order unity, then one can again expect $\sim 10\%$ value for the evaporated gas fraction. Alternatively, cluster mergers can lead to some gas loss. However, this process is episodic and is already taken into account in LSS hydro simulations.

We thank the referee, Alex Lazarian, for discussion and comments. This work has been supported by grants NNG04GM41G, NNX07AJ50G and DEFG0204ER54790.

REFERENCES

- Afshordi, N., Lin, Y.-T., Nagai, D., & Sanderson, A. J. R., 2006 astro-ph/0612700
 Chandran, B. D. G., & Cowley, S. C. 1998, Phys. Rev. Lett., 80, 3077
 Dolag, K., Vazza, F., Brunetti, G., Tormen, G. 2005, MNRAS, 364, 753
 Enßlin, T. A., Vogt, C. Pfrommer, C. 2005, in Proc. Magnetized Plasma in Galaxy Evolution (eds. Chyzy, K. T., Dettmar, R.-J., Otmainska-Mazur, K., Soida, M.), p.231; astro-ph/0501338
 Lazarian, A. 2006, ApJ, 645, L25
 Lazarian, A. 2007, ApJ, 660, L173
 Lesieur, M. 1990, Turbulence in Fluids: Stochastic and Numerical Modeling (2nd ed.; Dordrecht: Kluwer)
 Loeb, A. 2006, astro-ph/0606572
 Loeb, A. 2006, Notes for the SAAS-Fee Winter School (to be published by Springer-Verlag), astro-ph/0603360
 Malyshkin, L., & Kulsrud, R. 2001, ApJ, 549, 402
 Malyshkin, L. 2001, ApJ, 554, 561
 Narayan, R., & Medvedev, M. V. 2001, ApJ, 562, L129
 Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, ApJ, 490, 493
 Rechester, A. B., & Rosenbluth, M. N. 1978, Phys. Rev. Lett., 40, 38
 Spergel, D. N., et al. 2006, astro-ph/0603449
 Subramanian, K., Shukurov, A., Haugen, N. E. L. 2006, MNRAS, 366, 1437
 Vogt, C. & Enßlin, T. A. 2005, A&A, 434, 67
 Vogt, C. & Enßlin, T. A. 2003, A&A, 412, 373
 Vazza, F., Tormen, G., Cassano, R., Brunetti, G., Dolag, K. 2006, ApJ, 369, L14
 Voit, G. M. 2005, Reviews of Modern Physics, 77, 207




FIG. 1.— Fraction of the ICM gas evaporated from a cluster out of r_{vir} as a function of the temperature of the ICM gas. The solid curve is computed for $l_B = 1$ kpc and $c = 2$, the dotted is for $l_B = 10$ kpc and $c = 2$, and the dashed is for $l_B = 1$ kpc and $c = 8$.




FIG. 2.— Fraction of the evaporated gas as a function of l_B , the maximum correlation length the magnetic field. Here it runs from the observed value of ~ 1 kpc to the maximum possible scale, the cluster size, ~ 3 Mpc. The solid curve is computed for $T = 5$ keV and $c = 2$, the dotted is for $T = 15$ keV and $c = 2$, and the dashed is for $T = 5$ keV and $c = 8$.